

Effect of ursodeoxycholic acid on methionine adenosyltransferase activity and hepatic glutathione metabolism in rats

C M Rodríguez-Ortigosa, R N Cincu, S Sanz, F Ruiz, J Quiroga, J Prieto

Gut 2002;**50**:701–706

See end of article for authors' affiliations

Correspondence to:
Dr C M Rodríguez-Ortigosa,
Division of Hepatology and
Gene Therapy, School of
Medicine, University of
Navarra, Irunlarrea 1,
31080-Pamplona, Spain;
crodriguez@unav.es

Accepted for publication
17 July 2001

Background and aims: Both bile salts and glutathione participate in the generation of canalicular bile flow. In this work, we have investigated the effect of different bile salts on hepatic glutathione metabolism.

Methods: Using the isolated and perfused rat liver, we studied hepatic glutathione content, and metabolism and catabolism of this compound in livers perfused with taurocholate, ursodeoxycholate, or deoxycholate.

Results: We found that in livers perfused with ursodeoxycholate, levels of glutathione and the activity of methionine adenosyltransferase (an enzyme involved in glutathione biosynthesis) were significantly higher than in livers perfused with other bile salts. In ursodeoxycholate perfused livers, methionine adenosyltransferase showed a predominant tetrameric conformation which is the isoform with highest activity at physiological concentrations of substrate. In contrast, the dimeric form prevailed in livers perfused with taurocholate or deoxycholate. The hepatic activities of γ -glutamylcysteine synthetase and γ -glutamyltranspeptidase, enzymes involved, respectively, in biosynthetic and catabolic pathways of glutathione, were not modified by bile salts.

Conclusions: Ursodeoxycholate specifically enhanced methionine adenosyltransferase activity and hepatic glutathione levels. As glutathione is a defensive substance against oxidative cell damage, our observations provide an additional explanation for the known hepatoprotective effects of ursodeoxycholate.

Glutathione is a tripeptide involved in a diversity of hepatocellular metabolic pathways. It has been shown that glutathione is the main promoter of the bile salt independent fraction of canalicular flow.¹ In addition, glutathione is the principal intracellular defence of the liver against oxidative stress.² The biosynthetic pathway of this compound has two main characteristics: the availability of cysteine and synthesis of glutathione from this amino acid.³ Cysteine can be obtained from the diet, from methionine (through the transsulphuration pathway), or as a subproduct of glutathione catabolism.⁴

In the conversion of cysteine to glutathione, the limiting enzyme is γ -glutamylcysteine synthetase (γ -GCS) which is controlled by negative feedback from its end product glutathione⁵ whereas the formation of cysteine involves several reactions in the named methionine cycle. This cycle begins with incorporation of an ATP molecule to methionine to form S-adenosylmethionine (AdoMet) in an irreversible reaction catalysed by the enzyme methionine adenosyltransferase (MAT).⁶ In mammals, AdoMet is mainly used in transmethylation reactions⁷ that produce S-adenosylhomocysteine. This metabolite is converted to homocysteine which either rapidly enters the transsulphuration pathway to yield glutathione or it is used for resynthesis of methionine.⁸

It has been shown that in the liver there is a close relationship between glutathione content and MAT activity,^{9–11} suggesting that this enzyme plays an important role in hepatic glutathione metabolism. In the liver, MAT can be found in two isoforms, MAT I and MAT III, which are the tetramer and dimer forms, respectively, of the same polypeptide subunit.¹² These two isoforms have different kinetic properties and total MAT activity in the liver depends critically on the relative content of each.¹²

In the present study, we have investigated the effect of bile salts on hepatic glutathione metabolism in the isolated and perfused rat liver. Our data show that in contrast with other bile salts, ursodeoxycholic acid (UDC) specifically increases liver glutathione content by stimulating hepatic MAT activity. This paper provides evidence for a novel mechanism which contributes towards explaining the known hepatoprotective actions of UDC.

MATERIAL AND METHODS

Bile salts, ATP, methionine, glutathione reductase, bovine serum albumin, L- ∞ -aminobutyrate, pyruvate kinase, lactate dehydrogenase, and NADH were obtained from Sigma Chemical Co. (St Louis, Missouri, USA). [2-³H]ATP (25.5 Ci/mmol) was purchased from Amersham (Little Chalfont, UK). NYTRAN membranes were from Schleicher and Schuell (Keene, New Hampshire, USA). All other chemicals were of analytical reagent grade and were purchased from Merck (Darmstadt, Germany).

Livers from male Wistar rats (245–260 g body weight) were isolated and perfused in a recirculating anterograde way. The technique for liver isolation has been previously described.¹³ Briefly, after abdominal laparotomy the liver was exposed, and both the bile duct and portal vein were cannulated. The liver

Abbreviations: AdoMet, S-adenosylmethionine; DC, deoxycholate; γ -GCS, γ -glutamylcysteine synthetase; γ -GT, γ -glutamyltranspeptidase; iNOS, inducible nitric oxide synthase; MAT, methionine adenosyltransferase; MnSOD, Mn superoxide dismutase; TC, taurocholate; UDC, ursodeoxycholate; AST, aspartate aminotransferase; LDH, lactate dehydrogenase; SDS-PAGE, sodium dodecyl sulphate-polyacrylamide gel electrophoresis.

was perfused in situ with oxygenated Krebs-Ringer bicarbonate buffer to eliminate all blood components and the thoracic vena cava was then cannulated. After transferring the liver to an acrylic platform, it was placed in a thermostat controlled and humidified acrylic chamber, and perfused at 30 ml/min with 250 ml of Krebs-Ringer bicarbonate buffer containing: 116 mM NaCl, 4.7 mM KCl, 1.2 mM KH_2PO_4 , 1 mM MgSO_4 , 1.25 mM CaCl_2 , 25 mM NaHCO_3 , 5 mM glucose, 0.2 mM pyruvate, 2 mM lactate, and 1% bovine serum albumin. Immediately after the beginning of perfusion, a continuous infusion of different bile salts was initiated and maintained during the 120 minutes of the experiments. The perfusion medium was continuously gassed with 95% O_2 /5% CO_2 , pH was automatically maintained at 7.35–7.40 with an automatic autoburette (Radiometer, Copenhagen, Denmark), and perfusion pressure was monitored (Letica pressure transducer, Barcelona, Spain). Animals were treated humanely and study protocols were approved by the ethics committee of animal experimentation of our institution.

We established three experimental groups in which 40 $\mu\text{mol/h}$ of taurocholate (TC, $n=5$), ursodeoxycholate (UDC, $n=6$), or deoxycholate (DC, $n=4$) were infused for 120 minutes. The dose of 40 $\mu\text{mol/h}$ was chosen on the basis of the studies of Gores *et al* in which they found that this rate was suitable to maintain bile flow in the isolated rat liver.¹⁴ To test the effect of higher doses of TC, UDC, or DC, the study was repeated with 60 $\mu\text{mol/h}$ of these bile salts in three additional groups of animals ($n=4$ for each group). All bile produced in 10 minute intervals from minute 0 onwards was collected in preweighed polypropylene tubes containing 70 μl of 3% metaphosphoric acid to reduce spontaneous oxidation of glutathione in bile,¹⁵ and samples of hepatic effluent were obtained every 30 minutes. At the end of the experiments, several fragments of each liver were rapidly frozen and stored at -80°C until use.

The viability of the livers during the experiments was assessed by measuring aspartate aminotransferase (AST) and lactate dehydrogenase (LDH) in the effluent perfusate. Bile flow was calculated by gravimetry (assuming a bile density of 1), and total bile salts in bile and in the perfusate were quantified by the 3α -hydroxysteroid dehydrogenase method. Total glutathione in bile, hepatic effluent, and hepatic tissue was measured by the method of Tietze,¹⁶ with minor modifications. Perfusate levels of AST and LDH, and γ -glutamyltranspeptidase (γ -GT) activity in the livers were measured by conventional enzymatic methods, using a Cobas Fara analyser (Roche Diagnostics, Basel, Switzerland).

The hepatic biosynthetic pathway of glutathione was evaluated by measuring the activities of γ -GCS and MAT. For determination of hepatic γ -GCS activity, the liver tissue was homogenised (1:5) in 0.2 M Tris HCl buffer, pH 8.0, using an Ultraturrax tissue tearer (IKA Labor Technik, Staufen, Germany) at 22 000 rpm for 30 seconds on ice. Enzyme activity was measured following the method described by Seelig and Meister.¹⁷ Briefly, 50 μl of tissue homogenate were added to a 1 ml incubation mixture containing 0.1 M Tris HCl buffer, 150 mM KCl, 5 mM Na_2ATP , 2 mM phosphoenolpyruvate, 10 mM L-glutamate, 10 mM L- α -aminobutyrate, 20 mM MgCl_2 , 2 mM $\text{Na}_2\text{-EDTA}$, 0.2 mM NADH, 17 μg of pyruvate kinase, and 17 μg of LDH, and the decrease in absorbance at 340 nm was followed. A control assay was performed by omission of L- α -aminobutyrate in the presence of tissue sample. Enzyme activity was expressed as nmol NADH/min/mg protein. Protein was determined by the pyrogallol red-molybdate method¹⁸ using a Cobas Mira analyser (Roche Diagnostics, Basel, Switzerland).

Liver samples (about 1 g) used for determination of MAT activity were homogenised with a Potter-Elvehjem apparatus in four volumes of ice cold 10 mM Tris HCl (pH 7.5) containing 0.3 M sucrose with 0.1% β -mercaptoethanol, 1 mM benzamidine, and 0.1 mM phenylmethylsulphonyl fluoride. The




Figure 1 Bile flow (A) and biliary bile salt excretion (B) over the course of the experiments in livers continuously infused with 40 $\mu\text{mol/h}$ of taurocholate (TC; $n=5$), ursodeoxycholic acid (UDC; $n=6$), or deoxycholic acid (DC; $n=4$). Values are mean (SEM).

homogenate was centrifuged at 12 500 g for 20 minutes at 4°C , and the supernatant was centrifuged again at 105 000 g for 60 minutes at 4°C .¹⁹ The supernatant was used to quantify MAT activity as described previously.¹⁹ Also, a possible direct effect of bile salts on MAT activity was tested by using purified MAT III²⁰ in the presence of 30 or 500 μM of TC, UDC, or DC.

For northern blot analysis, total liver RNA was isolated by the guanidinium thiocyanate method.²¹ Aliquots (20 μg) of total RNA were denatured at 65°C for five minutes in 5% formaldehyde, 50% formamide, 8% glycerol, and then size fractionated by electrophoresis (20 mA, 15 hours) in a 0.9% agarose gel under denaturing conditions. RNAs were then blotted and fixed to NYTRAN membranes. Prehybridisation and hybridisation were carried out as described previously.²² MAT mRNA levels were measured using a 2.2 kb *EcoRI* cDNA fragment of rat liver MAT.²³ mRNA levels of inducible nitric oxide synthase (iNOS) were determined using a 0.8 kb *EcoRI/HindIII* fragment from the macrophage iNOS cDNA.²⁴ Mn superoxide dismutase (MnSOD) mRNA levels were determined using a 0.8 kb *EcoRI/HindIII* fragment from rat liver cDNA (generous gift of Dr D Massaro, Georgetown University, Washington DC, USA). Equal loading of the RNA gels was assessed by hybridisation with a probe specific for β -actin.²⁵

For determination of MAT protein levels, liver samples were homogenised in four volumes of ice cold 10 mM Tris/HCl (pH 7.5) containing 0.3 M sucrose, 0.1% β -mercaptoethanol, and protease inhibitors, as above. The homogenate was centrifuged for one hour at 100 000 g . Protein concentration was measured in the supernatants by the method of Bradford,²⁶ and equal amounts of protein (30 μg) were subjected to 10% sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE). Proteins were electrophoretically transferred to nitrocellulose membranes in a buffer containing 25 mM Tris, 192 mM glycine, and 20% (vol/vol) methanol. Membranes were blocked overnight with 1% low fat dry milk in Tris buffered saline containing 0.05% Tween 20. Immunodetection of MAT was performed using a rabbit anti-MAT antiserum²⁷ and




Figure 2 Aspartate aminotransferase (GOT), lactate dehydrogenase (LDH), and glucose levels, and variation in perfusion pressure during the experimental period (120 minutes) in livers continuously infused with 40 $\mu\text{mol/h}$ of taurocholate (TC; $n=5$), ursodeoxycholic acid (UDC; $n=6$), or deoxycholic acid (DC; $n=4$). Values are mean (SEM).

a horseradish peroxidase conjugated secondary antibody. Blots were developed by enhanced chemiluminescence according to the manufacturer's instructions (Dupont, Boston, Massachusetts, USA).

In order to evaluate the relative content of MAT I and MAT III, 200 mg of liver tissue were homogenised in 1 ml of the same buffer described above for determination of levels of MAT protein. Homogenates were centrifuged at 100 000 g for one hour and the supernatants (1 ml) were loaded onto 5 ml disposable polypropylene columns containing 1 ml bed




Figure 3 Hepatic content of total glutathione at the end of the experimental period (120 minutes) in livers continuously infused with 40 $\mu\text{mol/h}$ of taurocholate (TC; $n=5$), ursodeoxycholic acid (UDC; $n=6$), or deoxycholic acid (DC; $n=4$). Values are mean (SEM). Statistically significant differences are indicated.




Figure 4 (A) Biliary excretion of glutathione during the experimental period in livers perfused with 40 $\mu\text{mol/h}$ of taurocholate (TC; $n=5$), ursodeoxycholate (UDC; $n=6$), or deoxycholate (DC; $n=4$). Values are mean (SEM) of biliary excretion of glutathione in the previous 30 minutes. (B) Sinusoidal efflux of glutathione. Bars represent mean (SEM) glutathione release to the hepatic effluent during the experimental period (120 minutes) in the same groups as before.

volume of phenyl-Sepharose CL-4B equilibrated in buffer A (10 mM HEPES/Na (pH 7.5) with 1 mM EDTA and 10 mM MgSO_4). MAT I and MAT III were eluted from the phenyl-Sepharose column as previously described.²⁷ Briefly, MAT I was recovered in the flow through fraction (total volume of 4 ml) and after washing the column with 8 ml buffer A, MAT III was eluted with 4 ml of buffer A containing 50% dimethyl sulphoxide. Aliquots of the MAT I eluate (0.5 ml) were lyophilised and resuspended in 100 μl of 1 \times SDS-PAGE loading buffer. In parallel, 50 μl of the dimethyl sulphoxide fraction (MAT III) (total volume 4 ml) were diluted with 50 μl of the 2 \times loading buffer. Then, 40 μl of each preparation were analysed by SDS-PAGE and anti-MAT antibody immunoblotting, as described above.

Statistical comparisons between experimental groups were performed by the Mann-Whitney, Wilcoxon, and Kruskal-Wallis non-parametric tests.²⁸ Values are expressed as mean (SEM).

RESULTS

As observed in fig 1, bile flow and biliary bile salt excretion during the experimental period were similar in livers perfused with 40 $\mu\text{mol/h}$ of TC, UDC, or DC and concentrations of bile salts in the perfusate were also comparable in the three groups (TC 12.72 (4.03); UDC 14.39 (2.91); and DC 17.72 (3.27) μM ; NS). Likewise, liver viability, assessed by release into the effluent of AST, LDH, and glucose, was not significantly different between the groups (fig 2). Levels of oxygen uptake were also similar in all groups (TC 2.12 (0.09); UDC 2.24 (0.03); and DC 1.96 (0.08) $\mu\text{mol/min/g liver}$; NS), and were adequate to maintain sufficient oxygen consumption.²⁹ Finally, perfusion pressure at the beginning of perfusion was similar in all groups, and was maintained without significant changes during the experimental period (fig 2).

However, when we measured levels of glutathione in hepatic tissues collected at the end of the experiments, we




Figure 5 Hepatic activity of methionine adenosyltransferase (MAT) at the end of the experimental period (120 minutes) in livers perfused with 40 $\mu\text{mol/h}$ of taurocholate (TC; $n=5$), ursodeoxycholate (UDC; $n=6$), or deoxycholate (DC; $n=4$). Values are mean (SEM). Statistically significant differences are indicated.

observed a significant increase in glutathione content in livers perfused with UDC compared with TC or DC perfused livers (fig 3). As shown in fig 4, higher values of glutathione in UDC perfused livers were not due to differences in biliary excretion (fig 4A) or in the sinusoidal efflux of this compound (fig 4B). In addition, hepatic catabolism of glutathione, as estimated by γ -GT activity, was similar in the three experimental groups (36.1 (11.0), 37.5 (2.7), and 31.1 (8.3) mU/g liver, respectively, in TC, UDC, and DC groups; NS). To gain insight into the mechanisms responsible for the increased values of glutathione in livers perfused with UDC, we analysed the activity of the two main enzymes involved in glutathione synthesis, γ -GCS⁵ and MAT,⁸ in tissue samples collected at the end of the experiments. We observed no significant differences in hepatic γ -GCS activity among the groups (30.5 (5.5), 27.8 (5.0), and 21.9 (7.3) nmol/mg prot/min, in TC, UDC, and DC groups, respectively; NS) while MAT activity was significantly increased in livers perfused with UDC compared with the other groups ($p<0.05$) (fig 5). This increase in MAT activity was more pronounced when the livers were perfused with 60 $\mu\text{mol/h}$ of bile salts. In parallel, UDC perfused livers showed a higher hepatic content of glutathione with respect to the other groups (see table 1).

To test whether the observed rise in MAT activity was a consequence of increased MAT gene expression, levels of MAT mRNA and MAT protein were measured in all experimental groups. As shown in fig 6, no significant changes were observed between these groups with respect to both MAT mRNA and MAT protein levels, as determined by northern and western blots, respectively.

Table 1 Effect of bile salt infusion on liver methionine adenosyltransferase (MAT) activity and liver glutathione content

	Bile salt infused		
	TC	UDC	DC
MAT activity†			
Infusion rate			
40 $\mu\text{mol/h}$	150.7 (5.1)	284.9 (13.3)*	103.8 (9.5)
60 $\mu\text{mol/h}$	162.6 (15.8)	328.4 (81.1)*	109.8 (21.9)
Glutathione content‡			
Infusion rate			
40 $\mu\text{mol/h}$	6.22 (0.15)	7.53 (0.24)*	5.21 (0.53)
60 $\mu\text{mol/h}$	6.49 (1.3)	10.02 (1.0)*	5.52 (0.8)

†Values are expressed as pmol/min/mg protein.

‡Values are expressed as $\mu\text{mol/g}$ liver.

Livers were infused for 120 minutes with 40 or 60 $\mu\text{mol/h}$ of taurocholate (TC; $n=5$), ursodeoxycholate (UDC; $n=6$), or deoxycholate (DC; $n=4$).

* $p<0.05$ compared with the other groups with the same infusion rate.




Figure 6 mRNA expression and protein levels of methionine adenosyltransferase (MAT) in livers perfused for 120 minutes with 40 $\mu\text{mol/h}$ of taurocholate (TC; $n=5$), ursodeoxycholate (UDC; $n=6$), or deoxycholate (DC; $n=4$). Representative northern and western blots, respectively, are shown.

As both MAT activity and glutathione levels were affected by reactive nitrogen and oxygen species,^{30,31} we determined mRNA levels of iNOS and MnSOD, two enzymes implicated in the generation and metabolism of NO and O_2^- , respectively, as markers of oxidative and nitrosative stress in the different groups of livers. No significant changes were found in levels of these transcripts in the three experimental groups (data not shown), suggesting that variations in liver glutathione content and MAT activity induced by UDC are not due to an effect on oxygen or nitrogen free radical production.

Also, we investigated whether UDC could exert a direct effect on the activity of purified MAT in vitro. To verify this, the effect of increasing concentrations of TC, UDC, or DC was tested and compared with samples in which no bile salts were added. As can be seen in table 2, when the bile salt concentration increases, MAT activity decreases, independently of the bile salt used, although this effect was more evident in the samples in which DC was used. Nevertheless, at concentrations similar to those found in hepatocytes (that is, in the range 30–60 μM),³² no statistically significant differences were found between groups.

To analyse the mechanisms underlying the enhanced MAT activity in livers which received UDC infusion, we quantified the relative proportions of MAT I and MAT III in the three groups of livers by immunoblotting after separation of the two MAT isoenzymes using phenyl-Sepharose CL-4B chromatography.²⁷ As shown in fig 7A, the relative proportions of MAT I and MAT III were similar in TC and DC perfused livers. In contrast, in livers perfused with UDC, we observed an increase in the tetrameric isoform (MAT I) and a reduction in the dimeric isoform (MAT III), resulting in a marked decrease in the MAT III/MAT I ratio (4.5 (0.5) in the UDC group v 10 (4) and 14 (4) arbitrary units in TC and DC perfused livers, respectively; $p<0.05$). A representative immunoblot of MAT isoforms obtained from the cytosolic fraction of livers corresponding to the three experimental groups is shown in fig 7B.

Table 2 Effect of bile salts on methionine adenosyltransferase (MAT) activity in vitro

	Bile salt tested		
	TC	UDC	DC
Bile salt concn			
30 μM	119.8 (11.0)	115.6 (7.9)	101.6 (2.3)
500 μM	102.1 (6.7)	94.5 (1.5)	95.2 (2.7)
2 mM	95.2 (6.4)	79.5 (5.5)	60.9 (12.7)
6.6 mM	79.2 (7.6)	61.7 (2.4)	34.6 (8.1)*
10 mM	73.4 (5.9)	50.5 (5.2)	36.1 (7.9)*

MAT activity was measured in vitro in the presence of increasing concentrations of the three bile salts studied (taurocholate (TC), ursodeoxycholate (UDC), or deoxycholate (DC); $n=4-6$ measurements at every point).

Results are expressed as percentage with respect to those obtained in samples in which no bile salts were added.

* $p<0.05$ with respect to corresponding values in the TC group.




Figure 7 Methionine adenosyltransferase (MAT) III and MAT I isoenzymes obtained from the hepatic tissue of livers continuously infused with taurocholic acid (TC; n=5), ursodeoxycholic acid (UDC; n=6), or deoxycholic acid (DC; n=4). The isoenzymes were separated by phenyl-Sepharose chromatography and analysed by immunoblotting. (A) MAT III/MAT I ratio for the different experimental groups, obtained by scanning densitometry of the autoradiograms. Values are mean (SEM); *p<0.05 with respect to the other groups. (B) Representative immunoblots of MAT III (III) and MAT I (I).

DISCUSSION

Our data demonstrate that UDC specifically influences the metabolism of glutathione in the liver. We found that hepatic glutathione levels were higher in livers perfused with UDC than in those which received TC or DC. As biliary excretion of glutathione and release of glutathione to the perfusate were similar in UDC, TC, and DC groups, the increased stores of this compound in livers treated with UDC should be due to differences in glutathione synthesis or catabolism. Our data indicate that changes in glutathione catabolism are unlikely to contribute to this effect as the activity of γ -GT, the enzyme which mediates catabolism of glutathione at the canalicular level,³³ did not differ between groups. In contrast with this finding, other authors have reported that bile salts affect γ -GT activity; however, these data were generated with millimolar concentrations of bile salts in an in vitro model, and no information regarding the effect of UDC was reported.^{34,35}

The biosynthetic pathway of glutathione in the liver involves several enzymes. The most important are γ -GCS, which is the rate limiting enzyme in the conversion of cysteine to glutathione,⁵ and MAT, whose activity, as previously shown, is closely related to glutathione concentration. Our results showed that the activity of γ -GCS was similar in all groups. Concerning MAT, in the liver there are two different conformations of the enzyme, the tetrameric (MAT I) and dimeric (MAT III) forms, the former being the most active at physiological concentrations of substrates.²⁰ UDC favours the adoption of the tetrameric conformation and increases the MAT I/MAT III ratio. This effect of UDC could therefore explain the higher levels of glutathione observed in UDC perfused livers. In addition, it has been shown that high glutathione concentrations increases MAT activity.³¹⁻³⁶ Thus increased synthesis of glutathione by enhanced MAT activity could in turn stimulate this enzyme. Enhanced MAT activity occurred in the absence of changes in MAT protein or MAT mRNA levels, and without variations in oxidative or nitrosative status (as estimated by MnSOD and iNOS mRNA values) in the three experimental groups studied. Therefore, it seems that UDC might elevate hepatic glutathione stores by favouring the

tetrameric conformation of MAT, and both effects would enhance MAT activity. In agreement with our results, Mitsuyoshi *et al* have recently reported that UDC increases levels of glutathione in hepatocytes in vitro.³⁷ These authors however did not analyse in their model the activity of MAT, an enzyme which, as we have shown, appears to be importantly influenced by UDC in vivo.

Other mechanisms have been postulated to explain the beneficial effect of UDC, including protection of bile ducts against injury induced by hydrophobic bile salts³⁸ and stimulation of biliary secretion of toxic compounds.³⁹ Although these mechanisms cannot be excluded it should be mentioned that we did not find stimulation of biliary secretion of bile salts or glutathione in livers perfused with UDC, suggesting that activation of biliary secretory processes is not a relevant effect of UDC in our model.

Glutathione is an essential factor in the cell defence against oxidative insults,² and substrates for glutathione biosynthesis, such as AdoMet, have been shown to exert a protective effect against cholestasis and various forms of liver damage.⁴⁰⁻⁴⁵ As UDC stimulates the activity of MAT and enhances glutathione stores in the liver, our data reveal a new mechanism to account for the known hepatoprotective actions of this bile salt. The explanation for this unique effect of UDC could be related to the physicochemical and pharmacological properties of this bile salt that make it more hydrophilic than other bile salts tested in our experiments.³⁹

In summary, our results showed that UDC augments the concentration of glutathione in the liver by enhancing MAT activity. This effect is an additional mechanism to explain the ability of UDC to defend the liver against cholestatic damage and toxic insults.

ACKNOWLEDGEMENTS

We thank Dr D Massaro (Georgetown University, Washington DC, USA) for the generous gift of a 0.8 kb *EcoRI/HindIII* fragment from rat liver cDNA used to determine mRNA levels of Mn superoxide dismutase, and Drs M García, A González, and JI Monreal for their technical help. We also acknowledge the economic support of the Plan Nacional de Investigación y Desarrollo (ref SAF 98/0132), Europharma, and Knoll Laboratories.

Authors' affiliations

C M Rodríguez-Ortigosa, R N Cincin, S Sanz, F Ruiz, J Quiroga, J Prieto, Division of Hepatology and Gene Therapy, University of Navarra, Pamplona, 31080, Spain

REFERENCES

- Ballatori N, Truong AT. Glutathione as a primary osmotic driving force in hepatic bile formation. *Am J Physiol* 1992;**263**:G617-24.
- Jakoby WB. Detoxication: deconjugation and hydrolysis. In: Arias IM, Boyer JL, Fausto N, *et al*, eds. *The liver: biology and pathobiology*. New York: Raven Press, 1994;429-42.
- DeLeve LD, Kaplowitz N. Glutathione metabolism and its role in hepatotoxicity. *Pharmacol Ther* 1991;**52**:287-305.
- Wang W, Ballatori N. Endogenous glutathione conjugates: occurrence and biological functions. *Pharmacol Rev* 1998;**50**:335-56.
- Richman PG, Meister A. Regulation of gamma-glutamylcysteine synthetase by nonallosteric feedback inhibition by glutathione. *J Biol Chem* 1975;**250**:1422-6.
- Cantoni GL. S-Adenosylmethionine: A new intermediate formed enzymatically from L-methionine and adenosine-triphosphate. *J Biol Chem* 1953;**204**:403-16.
- Cantoni GL. Biochemical methylation: selected aspects. *Annu Rev Biochem* 1975;**44**:435-51.
- Mato JM, Alvarez L, Corrales FJ, *et al*. S-Adenosylmethionine and the liver. In: Arias IM, Boyer JL, Fausto N, *et al*, eds. *The liver: biology and pathobiology*. New York: Raven Press, 1994;461-70.
- Pajares MA, Durán C, Corrales F, *et al*. Modulation of rat liver S-adenosylmethionine synthetase activity by glutathione. *J Biol Chem* 1992;**267**:17598-605.
- Ruiz F, Corrales FJ, Miqueo C, *et al*. Nitric oxide inactivates rat hepatic methionine adenosyltransferase in vivo by S-nitrosylation. *Hepatology* 1998;**28**:1051-7.
- Corrales FJ, Ruiz F, Mato JM. In vivo regulation by glutathione of methionine adenosyltransferase S-nitrosylation in rat liver. *J Hepatol* 1999;**31**:887-94.

- 12 **Kotb M**, Mudd SH, Mato JM, *et al*. Consensus nomenclature for the mammalian methionine adenosyltransferase genes and gene products. *Trends Genet* 1997;**13**:51–2.
- 13 **Rodríguez-Ortigosa CM**, Vesperinas I, Qian C, *et al*. Taurocholate-stimulated leukotriene C₄ biosynthesis and leukotriene C₄-stimulated choleresis in isolated rat liver. *Gastroenterology* 1995;**108**:1793–801.
- 14 **Gores GJ**, Kost LJ, LaRusso NF. The isolated perfused rat liver: Conceptual and practical considerations. *Hepatology* 1986;**6**:511–17.
- 15 **Eberle D**, Clarke R, Kaplowitz N. Rapid oxidation *in vitro* of endogenous and exogenous glutathione in bile of rats. *J Biol Chem* 1981;**256**:2115–17.
- 16 **Tietze F**. Enzymic method for quantitative determination of nanogram amounts of total and oxidized glutathione: Applications to mammalian blood and other tissues. *Anal Biochem* 1969;**27**:502–22.
- 17 **Seelig GF**, Meister A. Glutathione biosynthesis; γ -Glutamylcysteine synthetase from rat kidney. *Methods Enzymol* 1985;**113**:379–90.
- 18 **Orsonneau JL**, Douet P, Massoubre C, *et al*. An improved pyrogallol red-molybdate method for determining total urinary protein. *Clin Chem* 1989;**35**:2233–6.
- 19 **Martín Duce A**, Ortiz P, Cabrero C, *et al*. S-Adenosyl-L-methionine synthetase and phospholipid methyltransferase are inhibited in human cirrhosis. *Hepatology* 1988;**8**:65–8.
- 20 **Cabrero C**, Puerta J, Alemany S. Purification and comparison of two forms of S-adenosyl-L-methionine synthetase from rat liver. *Eur J Biochem* 1987;**170**:299–304.
- 21 **Chomczynski P**, Sacchi N. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 1987;**162**:156–9.
- 22 **Thomas PS**. Hybridization of denatured RNA and small DNA fragments transferred to nitrocellulose. *Proc Natl Acad Sci USA* 1980;**77**:5201–5.
- 23 **Alvarez L**, Asunción M, Corrales F, *et al*. Analysis of the 5' non-coding region of rat liver S-adenosylmethionine synthetase mRNA and comparison of the Mr deduced from the cDNA sequence and the purified enzyme. *FEBS Lett* 1991;**290**:142–6.
- 24 **Xie QW**, Calaycay J, Mumford RA, *et al*. Cloning and characterization of inducible nitric oxide synthase from mouse macrophages. *Science* 1992;**256**:225–8.
- 25 **Levi A**, Elridge JD, Patterson BM. Molecular cloning of a gene sequence regulated by nerve growth factor. *Science* 1985;**229**:393–5.
- 26 **Bradford MM**. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding. *Anal Biochem* 1975;**72**:248–54.
- 27 **Mingorance J**, Álvarez L, Sánchez-Góngora E, *et al*. Site-directed mutagenesis of rat liver S-adenosylmethionine synthetase—Identification of a cysteine residue critical for the oligomeric state. *Biochem J* 1996;**315**:761–6.
- 28 **Miller RG**. *Simultaneous statistical inference*, 2nd edn. New York: Springer-Verlag, 1981.
- 29 **Sugano T**, Suda K, Shimada M, *et al*. Biochemical and ultrastructural evaluation of isolated rat liver systems perfused with a hemoglobin-free medium. *J Biochem* 1978;**83**:995–1007.
- 30 **Ávila MA**, Mingorance J, Martínez-Chantar ML, *et al*. Regulation of rat liver S-adenosylmethionine synthetase during septic shock: role of nitric oxide. *Hepatology* 1997;**25**:391–6.
- 31 **Sánchez-Góngora E**, Ruiz F, Wei A, *et al*. Interaction of liver methionine adenosyltransferase with hydroxyl radical. *FASEB J* 1997;**11**:1013–19.
- 32 **Hofmann AF**. Bile acid secretion, bile flow and biliary lipid secretion in humans. *Hepatology* 1990;**12**:17–25S.
- 33 **Speisky H**, Shackel N, Varghese G, *et al*. Role of hepatic γ -glutamyltransferase in the degradation of circulating glutathione: Studies in the intact guinea pig perfused liver. *Hepatology* 1990;**11**:843–9.
- 34 **Gardell SJ**, Tate SS. Effects of bile acids and their glycine conjugates on γ -glutamyl transpeptidase. *J Biol Chem* 1983;**258**:6198–201.
- 35 **Abbott WA**, Meister A. Modulation of gamma-glutamyl transpeptidase activity by bile acids. *J Biol Chem* 1983;**258**:6193–7.
- 36 **Corrales F**, Ochoa P, Rivas C, *et al*. Inhibition of glutathione synthesis in the liver leads to S-adenosyl-L-methionine synthetase reduction. *Hepatology* 1991;**14**:528–33.
- 37 **Mitsuyoshi H**, Nakashima T, Sumida Y, *et al*. Ursodeoxycholic acid protects hepatocytes against oxidative injury via induction of antioxidants. *Biochem Biophys Res Commun* 1999;**263**:537–42.
- 38 **Heuman DM**, Bajaj RS, Lin Q. Adsorption of mixtures of bile salt taurine conjugates to lecithin-cholesterol membranes: implications for bile salt toxicity and cytoprotection. *J Lipid Res* 1996;**37**:562–73.
- 39 **Beuers U**, Boyer JL, Paumgartner G. Ursodeoxycholic acid in cholestasis: Potential mechanisms of action and therapeutic applications. *Hepatology* 1998;**28**:1449–53.
- 40 **Bray GP**, Tredger JM, Williams R. S-adenosylmethionine protects against acetaminophen hepatotoxicity in two mouse models. *Hepatology* 1992;**15**:297–301.
- 41 **Osman E**, Owen JS, Burroughs AK. Review article: S-adenosyl-L-methionine—a new therapeutic agent in liver disease? *Aliment Pharmacol Ther* 1993;**7**:21–8.
- 42 **Stramentinoli G**, Gualano M, Ideo G. Protective role of S-adenosyl-L-methionine on liver injury induced by D-galactosamine in rats. *Biochem Pharmacol* 1978;**27**:1431–3.
- 43 **Boelsterli UA**, Rakhit G, Balazs T. Modulation by S-adenosyl-L-methionine of hepatic Na⁺, K⁺-ATPase, membrane fluidity, and bile flow in rats with ethinyl estradiol-induced cholestasis. *Hepatology* 1983;**3**:12–17.
- 44 **Gasso M**, Rubio M, Varela G, *et al*. Effects of S-adenosylmethionine on lipid peroxidation and liver fibrogenesis in carbon tetrachloride-induced cirrhosis. *J Hepatol* 1996;**25**:200–5.
- 45 **Cincu R**, Rodríguez-Ortigosa CM, Vesperinas I, *et al*. S-Adenosyl-L-methionine protects the liver against the cholestatic, cytotoxic, and vasoactive effects of leukotriene D₄: a study with isolated and perfused rat liver. *Hepatology* 1997;**26**:330–5.



Effect of ursodeoxycholic acid on methionine adenosyltransferase activity and hepatic glutathione metabolism in rats

C M Rodríguez-Ortigosa, R N Cincu, S Sanz, et al.

Gut 2002 50: 701-706

doi: 10.1136/gut.50.5.701

Updated information and services can be found at:

<http://gut.bmj.com/content/50/5/701.full.html>

These include:

References

This article cites 41 articles, 15 of which can be accessed free at:

<http://gut.bmj.com/content/50/5/701.full.html#ref-list-1>

Article cited in:

<http://gut.bmj.com/content/50/5/701.full.html#related-urls>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>